

MORENE

French Fries Production



CHARACTERISTICS

FLESH COLOUR	CREAM
SKIN COLOUR	WHITE
COOKING TYPE	BC
BERRIES	MANY
DRY MATTER	22.6%

PLANT POPULATIONS

TUBERS/ 50KGS	POPULATION/HA
500	29,500
600	31,000
700	33,000
800	35,000
950	37,000

These plant population guidelines are for irrigated crops in the absence of PCN.

FERTILISER ADVICE

Requires a moderate Nitrogen fertilisation.

SPECIAL ATTENTION POINTS

Before harvest haulm killing is recommended. The foliage should be mature. Mother tubers should be removed before storage.

1 = Susceptible, 9 = Resistant

BLACK LEG	4	
COMMON SCAB	7	
POWDERY SCAB	7	
DRY ROT	0.5	
SPRAING (TRV)	6.5	
FOLIAGE BLIGHT	6	
TUBER BLIGHT	7	
LITTLE POTATO DISORDER	8	
BRUISING	12	(1=Low, 20=Sensitive)
PCN RESISTANCE	RO 1	

MATURITY	5	(9=Early, 4=Late)
DORMANCY	7.5	(3=Short, 9=Long)
MATURE YIELD	8.5	(1=Low, 9=High)
TUBER NUMBER	7	(1=Low, 9=High)
SHAPE	OVAL	



Primary use:
FRENCH FRIES