

VIVALDI

Set Skin Salad



CHARACTERISTICS

| | |
|--------------|--------------|
| FLESH COLOUR | LIGHT YELLOW |
| SKIN COLOUR | YELLOW |
| COOKING TYPE | AB |
| BERRIES | FEW |
| DRY MATTER | 18.9% |

PLANT POPULATIONS

| TUBERS/ 50KGS | POPULATION/HA |
|---------------|---------------|
| 500 | 63,000 |
| 600 | 70,000 |
| 700 | 80,000 |
| 800 | 85,000 |
| 950 | 95,000 |

These plant population guidelines are for irrigated crops in the absence of PCN.

1 = Susceptible, 9 = Resistant

| | | |
|------------------------|-----|-----------------------|
| BLACK LEG | 7 | |
| COMMON SCAB | 5.5 | |
| POWDERY SCAB | 5 | |
| DRY ROT | 6.5 | |
| SPRAING (TRV) | 8 | |
| FOLIAGE BLIGHT | 5 | |
| TUBER BLIGHT | 8.5 | |
| LITTLE POTATO DISORDER | 7 | |
| BRUISING | 14 | (1=Low, 20=Sensitive) |
| PCN RESISTANCE | | |

FERTILISER ADVICE

Vivaldi is sensitive to Magnesium and Manganese deficiency. Field and foliar applications should be made. Give a start application of 80 kg nitrogen per ha.

SPECIAL ATTENTION POINTS

Treat as normal practice. Don't leave in the soil too long after haulm killing, because brightness of the skin will get poor.

| | | |
|--------------|------|-------------------|
| MATURITY | 7.5 | (1=Early, 9=Late) |
| DORMANCY | 6.5 | (1=Short, 9=Long) |
| MATURE YIELD | 8 | (1=Low, 9=High) |
| TUBER NUMBER | 8 | (1=Low, 9=High) |
| SHAPE | OVAL | |



Primary use:
Retail Fresh